2020 eCQM Flow – **CMS104v8: Discharged on Antithrombotic Therapy (STK-02)***

**This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.*

Measure Flow Diagram

2020 eCQM Flow – **CMS104v8: Discharged on Antithrombotic Therapy (STK-02)***

**This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.*

Measure Flow Diagram (Continued)

Sample Calculation

$$\text{Performance Rate} = \frac{\text{Numerator } (c = 50)}{\text{Denominator } (a = 100) - \text{Denominator Exclusions } (b1 + b2 = 20) - \text{Denominator Exceptions } (d1 + d2 = 20)} = 83\%$$

2020 eCQM Flow – **CMS104v8: Discharged on Antithrombotic Therapy (STK-02)***

**This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.*

Measure Flow Narrative

The measure flow diagram on the preceding pages illustrates the steps to determine the population criteria for this measure.

Measure Description	This measure assesses ischemic stroke patients prescribed or continuing to take antithrombotic therapy at hospital discharge
Initial Population	Start by identifying the initial population criteria as inpatient hospitalizations with: • patients age 18 and older discharged from inpatient care (non-elective admissions) • a principal diagnosis of ischemic or hemorrhagic stroke • a length of stay less than or equal to 120 days that ends during the measurement period
Denominator	The denominator criteria further constrains the initial population by inpatient hospitalizations for patients with a principal diagnosis of Ischemic stroke

Denominator Exclusions	The denominator exclusions criteria are used to identify a subset of the denominator population by excluding inpatient hospitalizations for patients with any of the following:
	• comfort measures documented
	• or a documented discharge disposition of:
	  – discharged to another hospital; or
	  – left against medical advice; or
	  – expired; or
	  – discharged to either a health care facility or home, for hospice care
	• or admitted for elective carotid intervention (implicit by inclusion of only non-elective hospitalizations)

2020 eCQM Flow – **CMS104v8: Discharged on Antithrombotic Therapy (STK-02)***

**This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.*

Measure Flow Narrative (Continued)

The measure flow diagram on the preceding pages illustrates the steps to determine the population criteria for this measure.

Numerator	The numerator criteria identify a subset of the denominator population (that did not meet the denominator exclusions criteria) by including inpatient hospitalizations for patients prescribed or continuing to take antithrombotic therapy at hospital discharge
Denominator Exceptions	The denominator exceptions criteria identify a subset of the denominator population (that did not meet the denominator exclusions criteria or numerator criteria) by excluding inpatient hospitalizations for patients who have either: • a documented reason for not prescribing antithrombotic therapy at discharge (either a qualifying medical or patient reason) • or received Ticagrelor as an antithrombotic therapy during the hospitalization