


2021 eCQM Flow – CMS72v9: Antithrombotic Therapy By End of Hospital Day 2 (STK-05)*

*This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.


Measure Flow Diagram



2021 eCQM Flow – CMS72v9: Antithrombotic Therapy By End of Hospital Day 2 (STK-05)*

*This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.


Measure Flow Diagram (Continued)



2021 eCQM Flow – CMS72v9: Antithrombotic Therapy By End of Hospital Day 2 (STK-05)*

*This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.

Measure Flow Diagram (Continued)



2021 eCQM Flow – CMS72v9: Antithrombotic Therapy By End of Hospital Day 2 (STK-05)*

**This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.*

Measure Flow Narrative

The measure flow diagram on the preceding pages illustrates the steps to determine the population criteria for this measure.

Measure Description	This measure assesses ischemic stroke patients administered antithrombotic therapy by the end of hospital day 2
Initial Population	Start by identifying the initial population criteria as inpatient hospitalizations with: <ul style="list-style-type: none">• patients age 18 and older discharged from inpatient care (non-elective admissions)• with a principal diagnosis of ischemic or hemorrhagic stroke• and a length of stay less than or equal to 120 days that ends during the measurement period
Denominator	The denominator criteria further constrains the initial population by inpatient hospitalizations for patients with a principal diagnosis of ischemic stroke
Denominator Exclusions	The denominator exclusions criteria identify a subset of the denominator population by excluding inpatient hospitalizations for patients with any of the following: <ul style="list-style-type: none">• a duration of stay less than 2 days• comfort measures documented day of or the day after arrival• intra-venous or intra-arterial Thrombolytic (t-PA) Therapy administered within 24 hours prior to arrival or anytime during hospitalization

2021 eCQM Flow – CMS72v9: Antithrombotic Therapy By End of Hospital Day 2 (STK-05)*

**This flow diagram represents an overview of population criteria requirements. Please refer to the eCQM measure specification for a complete list of definitions, direct reference codes, data or timing elements included in this measure and required for submission.*

Measure Flow Narrative (Continued)

The measure flow diagram on the preceding pages illustrates the steps to determine the population criteria for this measure.

Numerator	The numerator criteria identify a subset of the denominator population (that did not meet the denominator exclusions criteria) by including inpatient hospitalizations for patients who had antithrombotic therapy administered the day of or day after hospital arrival
Denominator Exceptions	<p>The denominator exceptions criteria identify a subset of the denominator population (that did not meet the denominator exclusions criteria or numerator criteria) by excluding inpatient hospitalizations where:</p> <ul style="list-style-type: none">• patients have a documented reason for not administering antithrombotic therapy the day of or day after hospital arrival• patients who receive pharmacological contraindications for antithrombotic therapy, Ticagrelor or Prasugrel, the day of or day after hospital arrival• patients have an INR greater than 3.5