NOTE: This flow diagram represents an overview of population criteria requirements. Refer to the eCQM specification for a complete list of data elements included in this measure and required for submission.


Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents

Percentage of patients 3 to 17 years of age who had an outpatient visit with a primary care physician (PCP) or obstetrician/gynecologist (OB/GYN) and who had evidence of the following during the measurement period.

- 1. Percentage of patients with height, weight, and body mass index (BMI) percentile documentation
- 2. Percentage of patients with counseling for nutrition
- 3. Percentage of patients with counseling for physical activity

This eCQM is a patient-based measure


Initial Population 1 3 to 17 years of age by the end of the Hospice measurement care for any part of period the measurement period Do not include in Yes Initial Population **Denominator Exclusions 1** Denominator Exclusions 1 No Eligible (10 patients) encounter with a primary care physician or an obstetrician/gynecologist Diagnosis during the of pregnancy measurement any time during the period measurement period Denominator 1 No Equals Initial Go to Population 1 Page 2 (100 patients)




Performance Rate 1 =	
Numerator 1 (a ¹ = 60 patients)	60
Denominator 1 (b1 = 100 patients) - Denominator Exclusions 1 (x1 = 10 patients)	= 90 = 67 %
Stratification 1 of Performance Rate 1 =	
Numerator 1 (a² = 30 patients)	= 30 = 33 %
Denominator 1 ($b^1 = 100$ patients) - Denominator Exclusions 1 ($x^1 = 10$ patients)	90 = 33 %
Stratification 2 of Performance Rate 1 =	
Numerator 1 (a³ = 30 patients)	30
Denominator 1 ($b^1 = 100$ patients) - Denominator Exclusions 1 ($x^1 = 10$ patients)	= = 33 %


Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents


Initial Population 2





Performance Rate 2 =			
Numerator 2 (a ⁴ = 40 patients)		40	= 44 %
Denominator 2 (b² = 100 patients) - Denominator Exclusions 2 (x² = 10 patients)		= 90 =	= 44 %
Stratification 1 of Performance Rate 2 =			
Numerator 2 (a ⁵ = 20 patients)		20	00.0
Denominator 2 (b² = 100 patients) - Denominator Exclusions 2 (x² = 10 patients)		90	= 22 %
Stratification 2 of Performance Rate 2 =			
Numerator 2 (a ⁶ = 20 patients)		20	





erformance Rate 3 =	
lumerator 3 ($a^7 = 40$ patients)	40
enominator 3 (b³ = 100 patients) - Denominator Exclusions 3 (x³ = 10 patients)	90 = 44 %
stratification 1 of Performance Rate 3 =	
umerator 3 (a ⁸ = 20 patients)	$=\frac{20}{90}=22\%$
enominator 3 (b³ = 100 patients) - Denominator Exclusions 3 (x³ = 10 patients)	= 90 = 22 %
tratification 2 of Performance Rate 3 =	
lumerator 3 (a ⁹ = 20 patients)	20

eCQM Identifier: CMS155v13

eCQM Title: Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents

NOTE: This flow diagram represents an overview of population criteria requirements. Refer to the eCQM specification for a complete list of data elements included in this measure and required for submission.

Description: Percentage of patients 3 to 17 years of age who had an outpatient visit with a primary care physician (PCP) or obstetrician/gynecologist (OB/GYN) and who had evidence of the following during the measurement period.

- 1. Percentage of patients with height, weight, and body mass index (BMI) percentile documentation.
- 2. Percentage of patients with counseling for nutrition.
- 3. Percentage of patients with counseling for physical activity.

This eCQM is a patient-based measure.

eCQM Flow Narrative

Initial Population 1

Start by identifying the Initial Population 1, which includes patients 3 to 17 years of age by the end of the measurement period with an eligible encounter with a primary care physician (PCP) or an obstetrician/gynecologist (OB/GYN) during the measurement period.

Denominator 1

Denominator 1 equals the Initial Population 1. In the sample calculation provided at the end of the eCQM flow, Denominator 1 is equal to 100 patients.

Denominator Exclusions 1

The Denominator Exclusions 1 criteria identify a subset of the Denominator 1 population by excluding patients who are in hospice care for any part of the measurement period and patients with a diagnosis of pregnancy any time during the measurement period. In the sample calculation provided at the end of the eCQM flow, the Denominator Exclusions 1 are equal to 10 patients.

Numerator 1

The Numerator 1 criteria identify a subset of the Denominator 1 population (that did not meet the Denominator Exclusions 1 criteria) by including patients whose height, weight and body mass index (BMI) percentile were recorded during the measurement period. In the sample calculation provided at the end of the eCQM flow, Numerator 1 is equal to 60 patients.

Stratification 1

There are two strata for reporting separately by age group. Stratification 1 includes patients 3 to 11 years of age at the end of the measurement period. In the sample calculation provided at the end of the eCQM flow, Stratification 1 is equal to 30 patients.

Stratification 2

There are two strata for reporting separately by age group. Stratification 2 includes patients 12 to 17 years of age at the end of the measurement period. In the sample calculation provided at the end of the eCQM flow, Stratification 2 is equal to 30 patients.

Sample Calculation: Population 1

A sample calculation for Population 1 is provided to help determine how the measure performance rate 1 is derived. The measure performance rate 1 is calculated by dividing the Numerator 1 (total equals 60 patients) by the difference between the Denominator 1 (total equals 100 patients) and the Denominator Exclusions 1 (total equals 10 patients), which is equal to a score of 67 percent. Stratification 1 of performance rate 1 is calculated by dividing the Numerator 1 of Stratification 1 (total equals 30 patients) by the difference between the Denominator 1 (total equals 100 patients) and the Denominator Exclusions 1 (total equals 10 patients), which is equal to a score of 33 percent. Stratification 2 of performance rate 1 is calculated by dividing the Numerator 1 of Stratification 2 (total equals 30 patients) by the difference between the Denominator 1 (total equals 100 patients) and the Denominator Exclusions 1 (total equals 10 patients), which is equal to a score of 33 percent.

Initial Population 2

Continue by identifying the Initial Population 2, which includes patients 3 to 17 years of age by the end of the measurement period with an eligible encounter with a primary care physician (PCP) or an obstetrician/gynecologist (OB/GYN) during the measurement period.

Denominator 2

Denominator 2 equals the Initial Population 2. In the sample calculation provided at the end of the eCQM flow, Denominator 2 is equal to 100 patients.

Denominator Exclusions 2

The Denominator Exclusions 2 criteria identify a subset of the Denominator 2 population by excluding patients who are in hospice care for any part of the measurement period and patients with a diagnosis of pregnancy any time during the measurement period. In the sample calculation provided at the end of the eCQM flow, the Denominator Exclusions 2 are equal to 10 patients.

Numerator 2

The Numerator 2 criteria identify a subset of the Denominator 2 population (that did not meet the Denominator Exclusions 2 criteria) by including patients who received nutrition counseling during the measurement period. In the sample calculation provided at the end of the eCQM flow, Numerator 2 is equal to 40 patients.

Stratification 1

There are two strata for reporting separately by age group. Stratification 1 includes patients 3 to 11 years of age at the end of the measurement period. In the sample calculation provided at the end of the eCQM flow, Stratification 1 is equal to 20 patients.

Stratification 2

There are two strata for reporting separately by age group. Stratification 2 includes patients 12 to 17 years of age at the end of the measurement period. In the sample calculation provided at the end of the eCQM flow, Stratification 2 is equal to 20 patients.

Sample Calculation: Population 2

A sample calculation for Population 2 is provided to help determine how the measure performance rate 2 is derived. The measure performance rate 2 is calculated by dividing the Numerator 2 (total equals 40 patients) by the difference between the Denominator 2 (total equals 100 patients) and the Denominator Exclusions 2 (total equals 10 patients), which is equal to a score of 44 percent. Stratification 1 of performance rate 2 is calculated by dividing the Numerator 2 of Stratification 1 (total equals 20 patients) by the difference between the Denominator 2 (total equals 100 patients) and the Denominator Exclusions 2 (total equals 10 patients), which is equal to a score of 22 percent. Stratification 2 of performance rate 2 is calculated by dividing the Numerator 2 of Stratification 2 (total equals 20 patients) by the difference between the Denominator 2 (total equals 100 patients) and the Denominator Exclusions 2 (total equals 10 patients), which is equal to a score of 22 percent.

Initial Population 3

Continue by identifying the Initial Population 3, which includes patients 3 to 17 years of age by the end of the measurement period with an eligible encounter with a primary care physician (PCP) or an obstetrician/gynecologist (OB/GYN) during the measurement period.

Denominator 3

Denominator 3 equals the Initial Population 3. In the sample calculation provided at the end of the eCQM flow, Denominator 3 is equal to 100 patients.

Denominator Exclusions 3

The Denominator Exclusions 3 criteria identify a subset of the Denominator 3 population by excluding patients who are in hospice care for any part of the measurement period and patients with a diagnosis of pregnancy any time during the measurement period. In the sample calculation provided at the end of the eCQM flow, the Denominator Exclusions 3 are equal to 10 patients.

Numerator 3

The Numerator 3 criteria identify a subset of the Denominator 3 population (that did not meet the Denominator Exclusions 3 criteria) by including patients who received physical activity counseling during the measurement period. In the sample calculation provided at the end of the eCQM flow, Numerator 3 is equal to 40 patients.

Stratification 1

There are two strata for reporting separately by age group. Stratification 1 includes patients 3 to 11 years of age at the end of the measurement period. In the sample calculation provided at the end of the eCQM flow, Stratification 1 is equal to 20 patients.

Stratification 2

There are two strata for reporting separately by age group. Stratification 2 includes patients 12 to 17 years of age at the end of the measurement period. In the sample calculation provided at the end of the eCQM flow, Stratification 2 is equal to 20 patients.

Sample Calculation: Population 3

A sample calculation for Population 3 is provided to help determine how the measure performance rate 3 is derived. The measure performance rate 3 is calculated by dividing the Numerator 3 (total equals 40 patients) by the difference between the Denominator 3 (total equals 100 patients) and the Denominator Exclusions 3 (total equals 10 patients), which is equal to a score of 44 percent. Stratification 1 of performance rate 3 is calculated by dividing the Numerator 3 of Stratification 1 (total equals 20 patients) by the difference between the Denominator 3 (total equals 100 patients) and the Denominator Exclusions 3 (total equals 10 patients), which is equal to a score of 22 percent. Stratification 2 of

performance rate 3 is calculated by dividing the Numerator 3 of Stratification 2 (total equals 20 patients) by the difference between the Denominator 3 (total equals 100 patients) and the Denominator Exclusions 3 (total equals 10 patients), which is equal to a score of 22 percent.